HEALTH MART ASPIRIN- aspirin tablet, film coated McKesson

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

McKesson Aspirin Drug Facts

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)* *nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

temporarily relieves

- headache
- muscle pain
- toothache
- menstrual pain
- pain and fever of colds
- minor pain of arthritis

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

if you have ever had an allergic reaction to any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- redness or swelling is present
- fever gets worse or lasts more than 3 days
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

These could be signs of a serious condition

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

• drink a full glass of water with each dose

- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours
- children under 12 years: consult a doctor

Other information

• store at 20°-25°C (68°-77°F)

Inactive ingredients

corn starch, dibasic calcium phosphate dihydrate, hypromellose, talc, triacetin

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to Bayer® Aspirin active ingredient Adult Aspirin Pain Reliever/Fever Reducer (NSAID) Headaches • Everyday aches and pains Safe pain relief plus lifesaving benefits Talk to your doctor or other healthcare provider before using this product for your heart. Gluten Free ACTUAL SIZE 300 Coated Tablets, 325 mg



| HEALTH MART AS | SPIRIN | | | | | |
|--|-----------|----------------------|--------------------|-------------------|----------------|----------|
| aspirin tablet, film coated | | | | | | |
| | | | | | | |
| Product Information | | | | | | |
| Product T ype | HUMAN | OTC DRUG | Item Code (Source) | | NDC:62011-0020 | |
| Route of Administration | ORAL | | | | | |
| | | | | | | |
| | • | | | | | |
| Active Ingredient/Activ | 0 | | | | | |
| Ingredient Name | | | | Basis of Strength | | Strength |
| ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E) | | | ASPIRIN | | 325 mg | |
| | | | | | | |
| | | | | | | |
| Inactive Ingredients | | | | | | |
| | | ıgredient Name | | | | Strength |
| STARCH, CORN (UNII: 08232 | NY3SJ) | | | | | |
| DIBASIC CALCIUM PHO SPH | | E (UNII: O7TSZ97GEP) |) | | | |
| HYPROMELLOSES (UNII: 3NZ | XW29V3WO) | | | | | |
| TALC (UNII: 7SEV7J4R1U) | | | | | | |
| TRIACETIN (UNII: XHX3C3X6 | 73) | | | | | |
| | | | | | | |
| | | | | | | |
| Product Characteristics | 5 | | | | | |
| Color | WHITE | Score | | | no score | |
| Shape | ROUND | Size | | | 14mm | |
| Flavor | | Imprint Co | ode | | Aspirin;L | |

| Contains | | | | | | | | |
|---------------------------|--|----------------------|--------------------|--|--|--|--|--|
| | | | | | | | | |
| Packaging | | | | | | | | |
| # Item Code | Package Description | Marketing Start Date | Marketing End Date | | | | | |
| 1 NDC:62011-0020-1 | 100 in 1 BOTTLE; Type 0: Not a Combination Product | 11/21/2011 | | | | | | |
| 2 NDC:62011-0020-3 | 300 in 1 BOTTLE; Type 0: Not a Combination Product | 11/21/2011 | | | | | | |
| B NDC:62011-0020-2 | 500 in 1 BOTTLE; Type 0: Not a Combination Product | 02/08/2017 | | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| Marketing Information | | | | | | | | |
| Marketing Categor | y Application Number or Monograph Citation | Marketing Start Date | Marketing End Date | | | | | |
| OTC monograph not fina | al part343 | 11/21/2011 | | | | | | |
| | 1 | | | | | | | |

Labeler - McKesson (177667227)

Revised: 8/2018

McKesson