DOK- docusate sodium capsule Major Pharmaceuticals

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredient (in each softgel)

Docusate Sodium 100 mg

Purpose

Stool softener laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in 12 to 72 hours

Warnings

Do not use

if you are presently taking mineral oil, unless told to do so by a doctor.

Ask a doctor before use if you have

- stomach pain
- nausea
- vomiting
- noticed a sudden change in bowel habits that last over 2 weeks

Stop use and ask a doctor if

- you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be a signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

• take only by mouth. Doses may be taken as a single daily dose or in divided doses.

adults and children 12 years and over	take 1 to 3 softgels daily.
children 2 to under 12 years of age	take 1 softgel daily
children under 2 years	ask a doctor

Other information

- each softgel contains: sodium 6 mg
- store at 25°C (77°F);excursions permitted between 15-30°C (59-86°F)

Inactive ingredients

edible ink, FD&C Red #40, FD&C Yellow #6, gelatin, glycerin, polyethylene glycol, propylene glycol*, purified water sorbitan, sorbitol

*contains one or more of these ingredients

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to the active ingredient in Colace® Regular Strength Stool Softener**

DOK^{тм} 100 mg

Docusate sodium

Stool Softener Laxative

For Use as a Stool Softener in Treating & Avoiding Constipation

Softgels

**This product is not manufactured or distributed bt Purdue Products L.P., distributor of Colace® Regular Strength Stool Softener.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

Distributed by: Major® Pharmaceuticals 17177 N Laurel Park Drive, Suite 233 Livonia, MI 48152

Product Label

NDC 0904-6457-60 Compare to the active ingredient in Colace ^a Regular Strength Stool Softener* DOCK 100 mg Docusate sodium Stool Softener Laxative For Use as a Stool Softener in Treating & Avoiding Constipation	TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.	Drug Facts	Active ingredient (in each softgel) Purpose Docusate sodium 100 mgStool softener laxative Uses	 Eleverse occasional consequence (insignating) generally produces bowel movement in 12 to 72 hours Warmings 	Do not use if you are presently taking mineral oil, unless told to do so by a doctor.	e if yo	stomach pain mausea womiting noticed a sudden change in bowel habits that lasts over 2 weeks	Drug Facts (continued under label)	This product is not manufactured or distributed by Purdue Products L.P., distributor of Colace® Regular Strength Stool Softener.	aceuticals Park Drive, Suite 233	Livonia, MI 48152 Re-Order No. 700370		PLD-E11 3	PHILE A
	Call 1-877-753-3935 Monday-Friday 9AM-5PM EST	Questions or comments?	Inactive ingredients edible ink, FD&C red #40, FD&C yellow #6, gelatin, glycerin, polyethylene glycol, propylene glycol*, purified water, sorbitan, sorbitol *may contain this ingredient	■ store at 25°C (77°F); excursions permitted between 15-30°C (59-86°F)	Other information each softael contains: sodium 6 ma	children under 2 years ask a doctor	children 2 to under 12 take 1 softgel daily years of age	adults and children take 1-3 softgels 12 years and over daily	 Directions take only by mouth. Doses may be taken as a single daily dose or in divided doses. 	right away.	professional before use. Keep out of reach of children. In case of overdose, and modicel help or profession Southal Contor	you need to use a laxative for more than 1 week	you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition.	Drug Facts (continued) Stop use and ask a doctor if
DOK docusate sodium capsule														
Product Information														
Product Type	HUMAN OTC DRUG			Iteı	Item Code (Source)					NDC:0904-6457				
Route of Administration ORAL														
Active Ingredient/Active Moi	ety													
Ingredient Name Basis o							s of S	Streng	th	Stre	ngth			
DOCUSATE SODIUM (UNII: F05Q2T2JA0) (DOCUSATE - UNII:M7P27195AG) DOCUSATE SODIUM 100 mg														
Inactive Ingredients]	Ing	redient Na	ıme								5	Streng	th
FD&C RED NO.40 (UNII: WZB9127XC														
FD&C YELLOW NO. 6 (UNII: H77VEIS	3A8)													
GELATIN (UNII: 2G86QN327L) SORBITAN (UNII: 6O92ICV9RU)														
POLYETHYLENE GLYCOL, UNSPEC	IFIED (UNI	: 3WJO0SD	W1A)										
			2.10 2000											

PROPYLENE GLYCOL (UNII: 6DC9Q167V3)

WATER (UNII: 059QF0KO0R)

SORBITOL (UNII: 506T60A25R)										
GLYCERIN (UNII: PDC6A3C0OX)										
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)										
Р	roduct Charact	teristics								
Color ORANGE Score					no score					
S	hape OVAL Size			12mm						
Fl	Flavor Imprint Code				P51;S77;SCU1;D2					
С	ontains									
Packaging										
#	Item Code		Package	Description	Marketing Start Date	Marketing End Date				
1	NDC:0904-6457- 60	100 in 1 B Product	OTTLE, PLASTIC;	Type 0: Not a Combination	02/28/2015					
2	NDC:0904-6457- 80	1000 in 1 Product	BOTTLE, PLASTIC	C; Type 0: Not a Combination	02/28/2015					
_										
Marketing Information										
Marketing Category			Application Nu	nber or Monograph Citation	Marketing Start Date	Marketing End Date				
0	TC MONOGRAPH N	OT FINAL	part334		02/28/2015					

Labeler - Major Pharmaceuticals (191427277)

Revised: 11/2019

Major Pharmaceuticals